



# CROP DISEASE SAMPLING GUIDE

Ascochyta blight

Canola blackleg

Septoria tritici blotch

Yellow spot

Septoria nodorum blotch

Net blotches

## Western Australia Sampling

Crop disease samples help us monitor the development of fungicide resistance in Australia

Thank you for supporting our research



Curtin University

# CROP DISEASE SAMPLING GUIDE

Ascochyta blight

Canola blackleg

Septoria tritici blotch

Yellow spot

Septoria nodorum blotch

Net blotches

1. Cut infected plant material with disease symptoms (stems, leaves or pods). Remove all uninfected parts to avoid moisture. We do not need whole plants.
2. Place samples into the provided yellow envelopes and seal. Samples from the same paddock (same variety and fungicide treatment) can be placed in the same yellow envelope, otherwise place samples in separate yellow envelopes.
3. Please indicate sample information (cultivar, location, treatment) on the yellow envelope or the information sheet provided.
4. Place your samples into the prepaid envelope along with the sample information sheet.
5. Place in the nearest mail box.

If you wish to sample a disease other than the one listed above please contact us at:

**Fungicide Resistance Group**

Centre for Crop and Disease Management

Email: [frg@curtin.edu.au](mailto:frg@curtin.edu.au)

Tel: +61 8 9266 1204

Web: [ccdm.com.au/research/fungicide-resistance-group](http://ccdm.com.au/research/fungicide-resistance-group)